

Convergence in capacity

Urban Cegrell

February 8, 2008

1 Introduction

The purpose of this paper is to study convergence of Monge-Ampère measures associated to sequences of plurisubharmonic functions defined on a hyperconvex subdomain Ω of $\mathbb{C}^n$.

This paper is an updated version of the preprint [C0].

The concept of convergence in capacity was introduced in [X], where it was proved that for any uniformly bounded sequence φ_j of plurisubharmonic functions that converges in $\mathbb{C}^n$ -capacity we have that $(dd^c\varphi_j)^n$ converges weak* to $(dd^c\varphi)^n, j \rightarrow +\infty$.

We generalize this result:

Theorem 1.1. *Assume $\mathcal{F} \ni u_0 \leq u_j \in \mathcal{F}$ and that u_j converges to u in $\mathbb{C}^n$ -capacity. Then $(dd^cu_j)^n$ converges weak* to $(dd^cu)^n, j \rightarrow +\infty$.*

We first recall some definitions. See [C1] and [C2] for details.

The class $\mathcal{F}$ consists of all plurisubharmonic functions φ on Ω such that there is a sequence $\varphi_j \in \mathcal{E}_0, \varphi_j \searrow \varphi, j \rightarrow +\infty$ and $\sup_j \int_{\Omega} (dd^c\varphi_j)^n < +\infty$, where $\mathcal{E}_0$ is the class of bounded plurisubharmonic functions ψ such that $\lim_{z \rightarrow \xi} \psi(z) = 0, \forall \xi \in \partial\Omega$ and $\int_{\Omega} (dd^c\psi)^n < +\infty$.

The following definition was introduced in [X]: A sequence $\varphi_j \in \mathcal{F}$ converges to φ in $\mathbb{C}^n$ -capacity if

$$\text{cap}(\{z \in k : |\varphi - \varphi_j| > \delta\}) \rightarrow 0, j \rightarrow +\infty \quad \forall k \subset\subset \Omega, \forall \delta > 0.$$

For $\omega \subset\subset \Omega, \text{cap}(\omega) = \int_{\Omega} (dd^c h_{\omega}^*)^n$ where h_{ω}^* is the smallest upper semicontinuous majorant of

$$h_{\omega}(z) = \sup\{\varphi(z); \varphi \in \mathcal{E}_0; \varphi|_{\omega} \leq -1\}.$$

Finally, we write $u_j \rightsquigarrow u, j \rightarrow +\infty$ if u_j converges weak* to $u, j \rightarrow +\infty$.

2 Proofs

Lemma 2.1. *Suppose μ is a positive measure on Ω which vanishes on all pluripolar sets and $\mu(\Omega) < +\infty$. If $u_0, u_j \in \mathcal{F}$, $u_0 \leq u_j \rightsquigarrow u$, $j \rightarrow +\infty$ and if $\int u_0 d\mu > -\infty$, then $\lim_{j \rightarrow +\infty} \int u_j d\mu = \int u d\mu$.*

Proof. Denote by dV the Lebesgue measure and choose $\tilde{u}_j \in \mathcal{E}_0 \cap C(\bar{\Omega})$, $\tilde{u}_j \geq u_j$ such that $\int_{\Omega} (\tilde{u}_j - u_j)(d\mu + dV) < \frac{1}{j}$. Then $\tilde{u} \rightsquigarrow u$, $\lim_{j \rightarrow +\infty} \int u_j d\mu - \int \tilde{u}_j d\mu = 0$ so it is enough to prove that

$$\lim_{j \rightarrow +\infty} \int \tilde{u}_j d\mu = \int u d\mu.$$

Thus we can assume $u_j \in \mathcal{E}_0 \cap C(\bar{\Omega})$.

By Theorem 6.3 in [C1] there is a $\psi \in \mathcal{E}_0$, $f \in L^1((dd^c \psi)^n)$ with

$$\mu = f(dd^c \psi)^n$$

so by lemma 5:2 in [C1], for every $p < +\infty$,

$$\lim_{j \rightarrow +\infty} \int u_j d\mu_p = \int u d\mu_p$$

there $\mu_p = \min(f, p)(dd^c \psi)^n$.

Now

$$\begin{aligned} \lim_{j \rightarrow +\infty} \int u_j d\mu &= \lim_{j \rightarrow +\infty} \int u_j d\mu_p + \\ &+ \lim_{j \rightarrow +\infty} \int u_j (f - \min(f, p))(dd^c \psi)^n \geq \\ &\geq \int u d\mu_p + \int u_0 (f - \min(f, p)) d\mu \\ &\rightarrow \int u d\mu, \quad p \rightarrow +\infty \end{aligned}$$

by monotone convergence. On the other hand, by Fatous lemma,

$$\limsup_{j \rightarrow +\infty} \int u_j d\mu \leq \int u d\mu$$

which gives the desired conclusion. □

Proof. Proof of the theorem.

We prove that

$$\lim_{j \rightarrow +\infty} \int h(dd^c u_j)^n = \int h(dd^c u)^n, \quad \forall h \in \mathcal{E}_0,$$

which is enough by Lemma 3:1 in [C1].

Suppose $\omega_1, \omega_2, \dots, \omega_{n-1} \in \mathcal{F}$, $h \in \mathcal{E}_0$. It follows from the assumption that $u_j \rightsquigarrow u$, $j \rightarrow +\infty$ so $\lim_j \int \omega_1 * dd^c u_j \wedge dd^c \omega_2 \wedge \dots \wedge dd^c h = \int \omega_1 dd^c u \wedge \dots \wedge dd^c h$ by the lemma.

Suppose now that

$$\lim_j \int \omega_1 (dd^c u_j)^p \wedge dd^c \omega_{p+1} \wedge \dots \wedge dd^c h = \int \omega_1 (dd^c u)^p \wedge \dots \wedge dd^c h.$$

for $1 \leq q \leq p \leq n-2$. We claim

$$\begin{aligned} & \lim_j \int \omega_1 (dd^c u_j)^{p+1} \wedge dd^c \omega_{p+2} \wedge \dots \wedge dd^c \omega_{n-1} \wedge dd^c h = \\ & = \int \omega_1 (dd^c u)^{p+1} \wedge dd^c \omega_{p+2} \wedge \dots \wedge dd^c \omega_{n-1} \wedge dd^c h. \end{aligned}$$

Given $\varepsilon > 0$ choose $k \subset\subset \Omega$ such that $\{z \in \Omega; h < -\varepsilon\} \subset k$ and then a subsequence u_{j_t} such that

$$\int_{\Omega} \left(dd^c \sum_{t=1}^{\infty} h_{\{z \in k; |u - u_{j_t}| > \varepsilon\}}^* \right)^n < 1$$

and denote by

$$h_N = \max \left(\sum_{t=N}^{\infty} h_{\{z \in k; |u - u_{j_t}| > \varepsilon\}}^*, -1 \right).$$

Then $h_N \rightarrow 0$, $N \rightarrow +\infty$ outside a pluripolar set and

$$\begin{aligned} & \int \omega_1 (dd^c u_{j_k})^{p+1} \wedge \dots \wedge dd^c h - \int \omega_1 (dd^c u_{j_k})^p dd^c u \wedge \dots \wedge dd^c h = \\ & = \int (u_{j_k} - u) (dd^c u_{j_k})^p \wedge dd^c \omega_1 \wedge \dots \wedge dd^c h = \\ & = \int -h_N (u_{j_k} - u) (dd^c u_{j_k})^p \wedge dd^c \omega_1 \wedge \dots \wedge dd^c h + \\ & + \int (1 + h_N) (u_{j_k} - u) (dd^c u_{j_k})^p \wedge dd^c \omega_1 \wedge \dots \wedge dd^c (h - h_{\varepsilon}) + \\ & + \int (1 + h_N) (u_{j_k} - u) (dd^c u_{j_k})^p \wedge dd^c \omega_1 \wedge \dots \wedge dd^c h_{\varepsilon} = I_{j_k} + II_{j_k} + III_{j_k}. \end{aligned}$$

where $h_{\varepsilon} = \max(h, -\varepsilon)$.

Thus

$$\begin{aligned}
|I_{j_k}| &\leq 2 \int (-h_N)(-u_0)(dd^c u_{j_k})^p \wedge \cdots \wedge dd^c h \leq \\
&\leq \int [-u_0 + \max(u_{0_1} - R)](dd^c u_{j_k})^p \wedge \cdots \wedge dd^c h + \\
&+ R \int -h_N (dd^c u_{j_k})^p \wedge \cdots \wedge dd^c h \\
&\rightarrow \int [-u_0 + \max(u_0, -R)](dd^c u)^p \wedge \cdots \wedge dd^c h + \\
&+ R \int (-h_N)(dd^c u)^p \wedge \cdots \wedge dd^c h, \quad j_k \rightarrow +\infty.
\end{aligned}$$

Now, the first term on the right hand side is small if R is large, the second is small if N is even larger.

$$\begin{aligned}
|III_{j_k}| &\leq \int -u_0 * (dd^c u_{j_k})^p \wedge dd^c \omega_1 \wedge \cdots \wedge dd^c h_\varepsilon \leq \\
&\leq \int -u_0 * (dd^c u_0)^p \wedge dd^c \omega_1 \wedge \cdots \wedge dd^c h_\varepsilon \leq \\
&\leq \int -h_\varepsilon (dd^c u_0)^{p+1} \wedge dd^c \omega_1 \wedge \cdots \wedge dd^c \omega_{n-1} \leq \\
&\varepsilon \int (dd^c u_0)^{p+1} \wedge dd^c \omega_1 \cdots \wedge dd^c \omega_{n-1} *
\end{aligned}$$

so it remains to estimate

$$\begin{aligned}
|II_{j_k}| &\leq 2\varepsilon \left(\int (dd^c u_{j_k})^p \wedge dd^c \omega_1 \cdots \wedge dd^c (h + h_\varepsilon) \right) \leq \\
&\leq 4\varepsilon \int (dd^c u_0)^p \wedge dd^c \omega_1 \cdots \wedge dd^c h.
\end{aligned}$$

which proves the claim.

Using the claim and repeating the chain of inequalities above, we conclude that

$$\begin{aligned}
&\int h(dd^c u_{j_k}^{**})^n - \int h(dd^c u)^n = \\
&\int (u_{j_k}^{**} * -u)(dd^c u_{j_k}^{**})^{n-1} \wedge h + \\
&\int u(dd^c u_{j_k}^{**})^{n-1} \wedge h - \int u(dd^c u)^{n-1} \wedge h \rightarrow 0, \quad j_k^{**} \rightarrow +\infty.
\end{aligned}$$

This proves the theorem, since we have proved that every subsequence of u_{j^*} contains a subsequence $u_{j_t^{**}}$ such that $(dd^c u_{j_t^{**}})^n$ converges weak* to $(dd^c u)^n$, $t \rightarrow +\infty$. □

References

- [C0] Cegrell, U., *Convergence in capacity*. Isaac Newton Institute for Mathematical Sciences. Preprint Series NI01046-NPD. Cambridge. 2001.
- [C1] Cegrell, U., *Pluricomplex energy*. Acta Math. **180:2** (1998), 187–217.
- [C2] Cegrell, U., *The general definition of the complex Monge-Ampère operator*. Ann. Inst. Fourier 54 (2004).
- [X] Xing, Y., *Continuity of the complex Monge-Ampère operator*. Proc. AMS., **124:2** (1996), 457–467.

February 8, 2008